

Derived equivalence of symmetric special biserial algebras

Mikhail Antipov
Saint-Petersburg University
hyperbor@list.ru

February 2, 2008

Abstract

We introduce Brauer complex of symmetric SB-algebra, and reformulate in terms of Brauer complex the so far known invariants of stable and derived equivalence of symmetric SB-algebras. In particular, the genus of Brauer complex turns out to be invariant under derived equivalence. We study transformations of Brauer complexes which preserve class of derived equivalence. Additionally, we establish a new invariant of derived equivalence of symmetric SB-algebras. As a consequence, symmetric SB-algebras with Brauer complex of genus 0 are classified.

Keywords: Brauer tree algebras, special biserial algebras, tilting complex

Contents

1	Introduction	1
2	The center $Z(\Lambda)$ and the multiplicities of A-cycles	2
3	Brauer complex	4
3.1	Definitions and constructions	4
3.2	Invariants of stable equivalence	4
4	Elementary tilting complexes	6
4.1	Definition of elementary tilting complex	6
4.2	Elementary transformations of Brauer complexes	7
4.3	Correspondence	8
5	Algebras of genus 0	10

1 Introduction

The present paper lies within a series of papers, devoted to classification of symmetric special biserial algebras up to derived equivalence (i.e., up to equivalence of derived categories). Recall that a symmetric SB-algebra Λ is uniquely determined by a pair $(\Gamma(\Lambda), f)$, where $\Gamma(\Lambda)$ is the

Brauer graph of Λ and $f : V(\Gamma(\Lambda)) \rightarrow \mathbb{N}$ maps vertices of $\Gamma(\Lambda)$ to their multiplicities (see, e.g., [1] and Proposition 3.9).

- We show that the multiset of multiplicities of vertices of $\Gamma(\Lambda)$ is invariant under derived equivalence (Proposition 2.1). In order to prove this, we determine the center $Z(\Lambda)$.
- In section 3 we introduce *Brauer CW-complex* $C(\Lambda)$ — a relevant tool for studying derived equivalence. Topologically, $C(\Lambda)$ is a sphere with handles. We reformulate in terms of $C(\Lambda)$ the basic notions related to algebra Λ and the invariants of stable equivalence, which appeared in [2]. In particular, the genus of $C(\Lambda)$ turns out to be invariant under stable equivalence. By a celebrated theorem of Rickard [7], these invariants are invariants of derived equivalence, too.
- We introduce *elementary tilting complexes* over symmetric special biserial algebras — a generalization of tilting complexes, which were treated in [3] (section 4). Equivalences of algebras, corresponding to elementary tilting complexes, can be reformulated in terms of 'elementary transformations' of Brauer CW-complexes of these algebras (Proposition 4.4). One sees that the algebra, which corresponds to the CW-complex obtained from $C(\Lambda)$ by an elementary transformation, is derived equivalent to Λ . Thus we obtain a direct graphic way of proving derived equivalence.
- In the last section we show that if the geometric realization of $C(\Lambda)$ is a sphere, then the invariants which we discuss in this paper determine Λ up to derived equivalence.

2 The center $Z(\Lambda)$ and the multiplicities of A -cycles

Let Λ be a symmetric SB-algebra over field K . Consider an extended quiver $Q_e = Q_e(\Lambda)$. Consider the partitions of its arrow set into A -cycles and into G -cycles (see[1]). Recall that A -cycles (and their multiplicities) correspond to the vertices of Brauer graph $\Gamma(\Lambda)$. We denote A -cycles by lower-case latine letters and denote vertices of $\Gamma(\Lambda)$ by the correspondent upper-case latine letters.

Let $\{c_1, c_2, \dots, c_k\}$ be the set of A -cycles. For each $i = 1, \dots, k$ consider a cyclic sequence $(\alpha_{i,1}, \alpha_{i,2}, \dots, \alpha_{i,l_i})$ of arrows of the cycle c_i . Let $f(c_1), f(c_2), \dots, f(c_k) \in \mathbb{N}$ denote the multiplicities of A -cycles. For each loop $\alpha = \alpha_{i,k}$ which is not formal, set

$$q_\alpha = (\alpha_{i,k+1}\alpha_{i,k+2} \dots, \alpha_{i,l_i} \dots \alpha_{i,k})^{f(c_i)-1} \alpha_{i,k+1}\alpha_{i,k+2} \dots, \alpha_{i,l_i} \dots \alpha_{i,k-1}.$$

Proposition 2.1. 1. *The center $Z(\Lambda)$ is generated as a vector space over K by 1 and by the elements of the following three forms:*

- Elements $m_{i,t} = (\alpha_{i,1}\alpha_{i,2} \dots, \alpha_{i,l_i})^t + (\alpha_{i,2}\alpha_{i,3} \dots, \alpha_{i,1})^t + \dots + (\alpha_{i,l_i}\alpha_{i,1} \dots, \alpha_{i,l_i-1})^t$ for all $i = 1, 2, \dots, k$ and $t = 1, \dots, f(c_i) - 1$.*
- Elements q_α for each non-formal loop α .*
- Elements $s_r = (\alpha_{i_r,1}\alpha_{i_r,2} \dots, \alpha_{i_r,l_{i_r}})^{f(c_{i_r})}$ for each vertex r of Q_e , where c_{i_r} is one of the two A -cycles, passing through r .*

2. $Z/(\text{Soc } Z) \cong K[x_1, x_2, \dots, x_k]/\langle \{x_i^{f(c_i)}, (x_i x_j)_{i \neq j}\} \rangle$, where $i, j \in 1, \dots, k$.
3. The multiset $(f(c_1), f(c_2), \dots, f(c_k))$ is invariant under derived equivalence.

Proof. **1.** Recall that the value of s_r doesn't depend on the choice of an A -cycle c_{i_r} and that the elements $s_1, s_2, \dots, s_n$ form a K -basis of $\text{Soc}(\Lambda)$ (see, e.g., [1]). Since Λ is a symmetric algebra, the socle $\text{Soc}(\Lambda)$ is contained in Z , so $s_r \in Z$. Moreover, for a non-formal loop α at vertex r and for the corresponding idempotent e_r and path $p \notin \{e_r, \alpha\}$ we get $e_r q_\alpha = q_\alpha = q_\alpha e_r$, $\alpha q_\alpha = s_r = q_\alpha \alpha$, $q_\alpha p = 0 = p q_\alpha$. Thus $q_\alpha \in Z$. Similarly, for all i, t, r we get $e_r m_{i,t} = m_{i,t} e_r$, since the summands in $m_{i,t}$ are circuits. Furthermore, for all l_1, l_2, t_1

$$\begin{aligned} (\alpha_{i,l_1} \alpha_{i,l_1+1} \dots, \alpha_{i,l_1-1})^{t_1} \alpha_{i,l_1} \alpha_{i,l_1+1} \dots, \alpha_{i,l_2} m_{i,t} = \\ (\alpha_{i,l_1} \alpha_{i,l_1+1} \dots, \alpha_{i,l_1-1})^{t+t_1} \alpha_{i,l_1} \alpha_{i,l_1+1} \dots, \alpha_{i,l_2} = \\ m_{i,t} (\alpha_{i,l_1} \alpha_{i,l_1+1} \dots, \alpha_{i,l_1-1})^{t_1} \alpha_{i,l_1} \alpha_{i,l_1+1} \dots, \alpha_{i,l_2} \end{aligned}$$

Since for the rest paths p (subpaths of other A -cycles) $m_{i,t} p = p m_{i,t} = 0$, we get $m_{i,t} \in Z$.

Each $z \in Z$ can be uniquely represented as

$$z = \sum_{j=1}^N a_j p_j + s, \quad (1)$$

where $0 \neq a_j \in K$, paths p_j are distinct nonzero paths in the quiver Q_e which are not contained in the socle, $s \in \text{Soc}(\Lambda)$.

By induction on the number of summands in the sum (1) we show, that z can be represented as a linear combination of elements $m_{i,t}$ and q_α . Fix $i \in \{1, \dots, N\}$ and write

$$p_i = \alpha_1 \alpha_2 \dots \alpha_m,$$

where $\alpha_1, \alpha_2, \dots, \alpha_m$ are consequent arrows of an A -cycle c_i . Let α_{m+1} be the next arrow of c_i . There are two cases:

Case 1: $\alpha_1 \alpha_2 \dots \alpha_m \alpha_{m+1} \notin \text{Soc}(\Lambda)$. In this case the path $\alpha_{m+1} \alpha_1 \alpha_2 \dots \alpha_m$ has coefficient a_i in the sum $\sum a_j \alpha_{m+1} p_j$. Since $z \alpha_{m+1} = \alpha_{m+1} z$, we obtain $\alpha_{m+1} = \alpha_1$, i.e. $p_i = (\alpha_{u,1} \alpha_{u,2} \dots, \alpha_{u,l_u})^t$ for some A -cycle c_u , $t < f(c_u)$. Moreover, the other summands of $m_{u,t}$ also have coefficient a_i in the sum (1). We see that the sum representing element $z - a_i m_{u,t} \in Z$ has less summands than the sum representing z , so the inductive hypothesis is applied.

Case 2: $\alpha_1 \alpha_2 \dots \alpha_m \alpha_{m+1} = s_l \in \text{Soc}(\Lambda)$ for some l . Consider an idempotent e_r such that $\alpha_m e_r \alpha_{m+1} \neq 0$. The expressions for $z e_r$ and $e_r z$ must contain p_i as a summand. Therefore p_i is a closed path. It follows that α_{m+1} is a loop and $p_i = q_{\alpha_{m+1}}$, and we apply the inductive hypothesis to $z - a_i p_i$.

2. Observe that $\text{Soc}(Z)$ is generated by the elements s_r and q_α , for all loops α which are not separate A -cycles (i.e., $\alpha q_\alpha = 0$). Moreover, $m_{i,t} m_{j,t_1} = \delta_{ij} m_{i,t+t_1}$ and $m_{i,t}^{f(c_i)} \in \text{Soc}(Z)$. These two observations imply the claim.

3. The claim follows directly from p.2, since $Z(\Lambda)$ is invariant under derived equivalence (see [6]). The maximal element $f(c_i)$ equals the maximal index of nilpotency of nilpotents in $Z/\text{Soc}(Z)$; the remaining proof is by induction. □

3 Brauer complex

3.1 Definitions and constructions

In this section we define a 2-dimensional CW-complex corresponding to a symmetric SB-algebra Λ . Associate with each G -cycle z of length k a k -gon F_z with an oriented border. The sides of F_z are labeled with the vertices of Q_e which lie on z (in the counter-clockwise order in the orientation of F_z). Consider a CW-complex $C = C(\Lambda)$ which is obtained from the resulting set of polygons by identifying oppositely oriented edges labeled by the same vertex. Since each vertex of Q_e belongs to exactly two G -cycles, C is an oriented manifold (without boundary).

Definition 3.1. CW-complex $C(\Lambda)$ is called *Brauer complex* of Λ .

Denote by $\Gamma = \Gamma(\Lambda)$ the Brauer graph of Λ . For a vertex $V \in V(\Gamma)$, consider a cyclic permutation π_V of half-edges, incident with V , which is defined by passing along the corresponding A -cycle. A 'picture' of a graph Γ on an oriented surface also determines, for any vertex of the graph, a cyclic permutation on the set of incident half-edges, which agrees with orientation. There exists an embedding i_Γ of Γ into an oriented surface M , which preserves the cyclic permutations (i_Γ and M are uniquely defined up to a homeomorphism). Note that we consider *strict* embeddings, i.e. such embeddings that each connectivity component of $M \setminus \Gamma$ is homeomorphic to an open disk). See [4] for the construction of embedding. It follows from the construction of embedding that the connectivity components of $M \setminus \Gamma$ correspond to the G -cycles of Λ . Now it is clear that M is a geometric realization of $C(\Lambda)$ and that the 1-skeleton S_M of $C(\Lambda)$ is isomorphic as a graph to Γ (we will refer to S_M as Γ). In particular, the vertices (edges) of $C(\Lambda)$ are in one-to-one correspondence with the A -cycles (resp., vertices) of Q_e . It is to be mentioned that the arrows of Q_e are in one-to-one correspondence with the angles of the 2-dimensional faces of $C(\Lambda)$.

Definition 3.2. *Perimeter* of a 2-dimensional face of $C(\Lambda)$ is the number of its edges, taking multiplicities into account (i.e., perimeter is the length of the corresponding G -cycles).

3.2 Invariants of stable equivalence

Observe that $C(\Lambda)$ is an oriented surface. The following statement holds since the Euler characteristic of an oriented surface is even.

Proposition 3.3. *If in the extended quiver Q_e of Λ the number of A -cycles is k , the number of G -cycles is g and the number of vertices is n , then $k + g - n$ is even.*

Remark 3.4. This statement was proved in [2] without topological arguments (Lemma 3.2).

Definition 3.5. The value $k + g - n$ is called the *genus* of Λ (and of $C(\Lambda)$).

In [2] it is proved that the multiset of lengths of G -cycles, as well as the number of A -cycles, is invariant under stable equivalence. By Rickard's Theorem, the derived equivalence of self-injective algebras implies stable equivalence (See [7]). The number of isomorphism classes of simple modules (i.e., the number of vertices of Q_e) is also stable invariant (See [5]). Therefore we get


Proposition 3.6. *The multiset of perimeters of faces, the number of vertices and the genus of $C(\Lambda)$ are invariant under derived equivalence.*

It was shown in [2] that the free rank of the Grothendieck group of the stable category $\text{stmod-}\Lambda$ equals $n - k$ if and only if $\Gamma(\Lambda)$ is not bipartite. Therefore, we have

Proposition 3.7. *Derived (stable) equivalence preserves the property of the Brauer graph to be bipartite.*

It should be mentioned that for algebras of genus 0 this invariant gives nothing new, since an embedded into a sphere graph is bipartite if and only if the perimeters of all its faces are even. But there are algebras of genus 1, the derived categories of which are not distinguished by the previously discussed invariants, but which are not equivalent by Proposition 3.7.

Example 3.8. Consider the following symmetric SB-algebras Λ_1 and Λ_2 :



The quiver Q_1 of Λ_1 consists of vertices 1, 2, 3 and arrows

$$\alpha, \delta : 1 \rightarrow 2, \beta, \varepsilon : 2 \rightarrow 3, \gamma, \eta : 3 \rightarrow 1.$$

Ideal I_1 of relations of Λ_1 is generated by the elements


$$\alpha\beta, \beta\gamma, \gamma\delta, \delta\varepsilon, \varepsilon\eta, \eta\alpha, \alpha\varepsilon\gamma - \delta\beta\eta, \varepsilon\gamma\alpha - \beta\eta\delta, \gamma\alpha\varepsilon - \eta\delta\beta.$$

The quiver Q_2 of Λ_2 consists of vertices 1, 2, 3 and arrows

$$\alpha_1 : 1 \rightarrow 2, \beta_1 : 2 \rightarrow 3, \gamma_1 : 3 \rightarrow 1, \delta_1 : 1 \rightarrow 3, \varepsilon_1 : 3 \rightarrow 2, \eta_1 : 2 \rightarrow 1.$$

Ideal I_2 of relations of Λ_2 is generated by the elements

$$\begin{aligned} \alpha_1\beta_1, \beta_1\gamma_1, \gamma_1\delta_1, \delta_1\varepsilon_1, \varepsilon_1\eta_1, \eta_1\alpha_1, \alpha_1\eta_1\delta_1\gamma_1 - \delta_1\gamma_1\alpha_1\eta_1, \\ \beta_1\varepsilon_1 - \eta_1\delta_1\gamma_1\alpha_1, \varepsilon_1\beta_1 - \gamma_1\alpha_1\eta_1\delta_1. \end{aligned}$$



It is easy to see that Λ_1 and Λ_2 are algebras with 3 simple modules, with one G -cycle of length 6 ($(\alpha\beta\gamma\delta\varepsilon\eta)$ and $(\alpha_1\beta_1\gamma_1\delta_1\varepsilon_1\eta_1)$, respectively) and with 2 A -cycles of multiplicities 1 ($c_1^1 = (\alpha\varepsilon\gamma)$, $c_2^1 = (\delta\beta\eta)$ and $c_1^2 = (\varepsilon_1\beta_1)$, $c_2^2 = (\gamma_1\alpha_1\eta_1\delta_1)$). In particular, Λ_1 and Λ_2 have genus 1. But $\Gamma(\Lambda_1)$ is bipartite (it consists of 2 vertices, connected by 3 edges) whereas $\Gamma(\Lambda_2)$ is not (the edge, corresponding the vertex 1 of Q_2 is a loop). Therefore, Λ_1 and Λ_2 are not derived equivalent.

Despite existence of an 'additional' invariant, the invariants and equivalences which are discussed in this paper are not enough to classify algebras of positive genus, in contrast to the 'spherical' case, which is treated in section 5 (see also example 4.7).

Proposition 3.9. *Correspondence $\Lambda \mapsto C(\Lambda)$ gives a bijection from the set of (pairly non-isomorphic) indecomposable symmetric SB-algebras to the set of (pairly non-isomorphic) pairs (C, f) , where*

1. C is a CW-complex homeomorphic to 2-dimensional oriented manifold with fixed orientation;

2. f is an arbitrary map from the 0-skeleton of C to $\mathbb{N}$.

Proof. It remains to show that a Brauer complex uniquely determines a symmetric SB-algebra. It follows from the fact the 1-skeleton of Brauer complex has a structure of Brauer graph, which uniquely determines a symmetric SB-algebra (see [1])¹. $\square$

4 Elementary tilting complexes

4.1 Definition of elementary tilting complex

Fix an edge i of C (equivalently, fix a vertex i in quiver Q_e), and suppose that there are other edges in C . We distinguish three cases.

1. i is a leaf of Γ . Equivalently, in the quiver Q_e there is a loop α_i at vertex i and this loop is an A -cycle (i.e., it annihilates all other arrows of Q_e).
2. i is a loop, which bounds some face of C . Equivalently, in the quiver Q_e there is a loop α_i at vertex i and this loop is a G -cycle. In this case there is a unique A -cycle passing through i (this cycle contains at least 3 arrows, one of which is α_i).
3. For $r = 1, 2$ the end $C_{i,r}$ of the edge i is incident with an edge $i_r \neq i$, such that $\pi_{C_{i,r}}(i_r) = i$. We permit $i_1 = i_2$ and we permit i to be a loop (i.e., $C_{i,1} = C_{i,2}$). Equivalently, there is no loop at vertex i of Q_e , i.e. the vertices i_1, i_2 which precede i on both A -cycles passing through A ($c_{i,1}$ and $c_{i,2}$) are different from i .

In each of these cases, to the edge i we put in correspondence a complex T_i as follows. For a vertex $j \in V(Q_e)$ we denote by P_j the indecomposable left projective Λ -module, which corresponds to j . For $i \neq j$, denote by T_{ij} the complex $\cdots \rightarrow 0 \rightarrow P_j \rightarrow 0 \rightarrow \cdots$ concentrated in degree 0. If i is a leaf of Γ , define complex T_{ii} by

$$T_{ii} : \cdots \rightarrow 0 \rightarrow P_j \xrightarrow{\beta_i} P_i \rightarrow 0 \rightarrow \cdots$$

where $j \in V(Q_e)$, $j \neq i$ is the vertex preceding vertex i on the (unique) G -cycle, which contains i ; $\beta_i \neq \alpha_i$ is the arrow preceding α_i on the same G -cycle.

If i is a loop which bounds some face of C , define T_{ii} by

$$T_{ii} : \cdots \rightarrow 0 \rightarrow P_j \bigoplus P_j \xrightarrow{(\beta_i, \beta_i \alpha_i)} P_i \rightarrow 0 \rightarrow \cdots$$

where $j \in V(Q_e)$, $j \neq i$ is the vertex preceding vertex i on the (unique) A -cycle, which contains i ; $\beta_i \neq \alpha_i$ is the arrow preceding α_i on the same A -cycle.

Otherwise, define T_{ii} by

$$T_{ii} : \cdots \rightarrow 0 \rightarrow P_{i_1} \bigoplus P_{i_2} \xrightarrow{(\beta_i^1, \beta_i^2)} P_i \rightarrow 0 \rightarrow \cdots$$

where i_1, i_2 are the vertices preceding i on the A -cycles $c_{i,1}$ and $c_{i,2}$, respectively; β_i^1, β_i^2 are the respective arrows preceding α_i . Finally, set $T_i = \bigoplus_{j=1}^n T_{ij}$.

¹ In [1] it was shown that a symmetric SB-algebra is uniquely determined by the (labeled) Brauer graph and certain parameters. It can be easily shown that these parameters are excessive and can be eliminated.

Proposition 4.1. T_i is a tilting complex over Λ .

Proof. We verify that T_i satisfies the two conditions from the definition of tilting complex. In the definition of T_i we distinguished three cases. We show verification only for the third case, the other cases are treated in the same way.

First, we must verify that $D^b(\Lambda) = \text{Add}(T_i)$, where $\text{Add}(T_i)$ is the smallest triangulated subcategory, which contains all direct summands of object T_i . It is enough to verify that all objects of the form $0 \rightarrow P_j \rightarrow 0$ belong to $\text{Add}(T_i)$. For $i \neq j$ this is by definition of T_i . For $i = j$ it is easy to see that $P_i[-1]$ is the third term of the triangle, which corresponds to the natural embedding of $T_{ii_1} \oplus T_{ii_2}$ into T_{ii} . It follows that T_i satisfies the first condition.

Now we verify that $\text{Hom}_{D^b(\Lambda)}(T_i, T_i[r]) = 0$ for $r \in \mathbb{Z} \setminus 0$. It is enough to prove that for each $j \in V(Q_e)$ $\text{Hom}_{D^b(\Lambda)}(T_{ii}, T_{ij}[-1]) = \text{Hom}_{D^b(\Lambda)}(T_{ij}[-1], T_{ii}) = 0$. Each morphism from T_{ij} to T_{ii} is determined by a morphism $f : P_j \rightarrow P_i$, where f is a multiplication by a linear combination of paths with starting point j and endpoint i . Each of these paths ends either with β_i^1 or with β_i^2 . Therefore f factors through $(\beta_i^1, \beta_i^2) : P_{i_1} \oplus P_{i_2} \rightarrow P_i$. It follows that f is homotopic to zero. Similarly, each morphism from T_{ii} to T_{ij} is determined by a morphism $f : P_i \rightarrow P_j$, where f is a multiplication by a linear combination S of paths with starting point i and endpoint j . Suppose that S has nonzero summands. Since $i \neq j$, the underlying paths are not maximal. Multiplying S by β_i^1 or by β_i^2 from the left, we again get a nontrivial sum of linearly independent summands. This contradicts the definition of morphism of complexes. Therefore $f = 0$ and $\text{Hom}_{D^b(\Lambda)}(T_{ii}, T_{ij}[-1]) = 0$. $\square$

4.2 Elementary transformations of Brauer complexes

Now we define *elementary transformations* of Brauer complexes. We will prove below that in terms of algebras, an elementary transformation puts an algebra Λ to the endomorphism algebra of one of the above defined tilting complexes over Λ . We fix convention that under elementary transformation the vertices are fixed, the configuration of edges (labeled with vertices of a quiver) — and therefore the configuration of faces (labeled with G -cycles) — is changed. In other words, we identify the edges (and faces) by their labels, not by the vertices incident to them. The pictures below illustrate the simplest cases, in general they can be quite different.

Definition 4.2. Let C be a Brauer complex, let Q_e be the corresponding extended quiver. Let $a \in E(C)$, $V \in V(C)$, let F be a face of C . Permutations $\text{Next}_F : V(C) \rightarrow V(C)$ and $E(C) \rightarrow E(C)$ are induced by the counter-clockwise order of vertices and edges in the orientation of F . Recall that π_V denotes the permutation of half-edges incident with vertex $V \in V(C)$, which is defined by passing along the corresponding A -cycle v . By abuse of language, we will name half-edges after correspondent edges. Thus by abuse of language for a loop a both situations $\pi_V(a) = a$ and $\pi_V(a) \neq a$ can happen. However, from the context it will always be clear which half-edge is meant.

4.2.1 Transformation of type 1: shift of a leaf

Let $V \in V(C)$ be a dangling vertex. Suppose that the edge (the face) incident with V is labeled by a (resp., by F). Let V_1 be the second vertex incident with a . Put $V_2 = \text{Next}_F(V_1)$, $a_1 = \text{Next}_F(a)$. Now shift edge a , so that a becomes incident with V and V_2 and $a = \text{Next}_F(a_1)$.




Figure 1: Shift of a leaf

4.2.2 Transformation of type 2: shift of a loop

Let a be a loop at vertex V_1 , bounding some face F_1 . Let F_2 be the second face, incident with a , put $V_2 = \text{Next}_{F_2}(V_1)$, $a_1 = \text{Next}_{F_2}(a)$. Replace loop a with a loop at vertex V_2 , which lies inside F_2 after a_1 . Note that F_1 is again bounded by a loop, which separates it from F_2 .




Figure 2: Shift of a loop

4.2.3 Transformation of type 3: the general case

Let a be an edge. Suppose that the vertices (faces) incident with a are labeled by V_1, V_2 (resp., by F_1 and F_2 ; we permit $F_1 = F_2$). For $i = 1, 2$ put $V_i' = \text{Next}_{F_i}^{-1}(V_i)$, $a_i = \text{Next}_{F_i}^{-1}(a)$. Shift a so that it becomes incident with V_1' and V_2' , separates F_1 from F_2 and lies after a_i on the new boundary of F_{3-i} .




Figure 3: The general case

Definition 4.3. We call the transformations of types 1-3 *tilting transformations*. The resulting complex is denoted by $C(a)$.

4.3 Correspondence

Proposition 4.4. Let Λ be an SB-algebra, $C = C(\Lambda)$, $a \in E(C)$. Let T_a be the tilting complex which corresponds to a . Then $\text{End}_{D^b(\Lambda)} T_a$ is a symmetric SB-algebra with Brauer complex $C(a)$

(C and $C(a)$ have the same multiplicities of vertices).

Proof. Denote by Q_e the extended quiver of Λ . By Rickard's theorem [6], $\Lambda_a = \text{End}_{D^b(\Lambda)} T_a$ is derived equivalent to Λ . Since Λ is a symmetric algebra, Λ_a is a symmetric algebra, too. By Pogorjaly's result, an algebra, which is stable equivalent to an SB-algebra, is an SB-algebra, too [5]. Therefore, by another Rickard's theorem [7] Λ_a is an SB-algebra. Let $e = \sum_{i=1}^n e_i$ be the decomposition of unity of Λ_a , which corresponds to the decomposition $T_a = \bigoplus_{i=1}^n T_{ai}$. Since the number of simple modules is invariant under derived equivalence, Λ_a is an algebra with n simple modules and therefore $\{e_i\}$ is a set of primitive orthogonal idempotents. Set $f_a = 1 - e_a \in \Lambda$, and denote $\Lambda_{-a} = f_a \Lambda f_a$. Since for $i \neq a$ the complexes T_{ai} are concentrated in degree 0, we have $\Lambda_{-a} = \text{End}_{\Lambda} \bigoplus_{i \neq a} P_i = \text{End}_{D^b(\Lambda)} \bigoplus_{i \neq a} T_{ai}$. Consider Brauer complex C_{-a} , obtained from C by deletion of an edge a (if a is a leaf, we delete it with the incident dangling vertex). The marks on the remaining vertices are preserved. We need the following lemma.

Lemma 4.5. *The symmetric SB-algebra which corresponds to C_{-a} is isomorphic to Λ_{-a} .*

Proof. We consider the case when $C(a)$ is obtained from C by a transformation of type 3 (i.e., a is not a loop which bounds a face and not a leaf). The other cases are treated in the same way. Denote the arrows of Q incident with a by $\alpha, \beta, \gamma, \delta$, so that $\alpha\beta \neq 0$ and $\gamma\delta \neq 0$. The elements of Λ_{-a} are linear combinations of paths whose starting points and endpoints differ from a . It is clear that Λ_{-a} is generated as algebra by idempotents e_i , where $i \neq a$, by arrows of Q different from $\alpha, \beta, \gamma, \delta$ and by the elements $\alpha\beta, \gamma\delta$. Observe that in terms of quivers Λ_{-a} can be obtained from Λ in the following way: the arrows α and β , lying on a common A -cycle, are replaced with an arrow $\alpha\beta$ on the same A -cycle (respectively, the arrows γ and δ are replaced with an arrow $\gamma\delta$). This implies the claim. $\square$

We return to the proof of proposition 4.4. Observe that the symmetric SB-algebra which corresponds to $C(a)_{-a} = C_{-a}$ is isomorphic to Λ_{-a} . To obtain the Brauer complex of Λ_a from C_{-a} we need to add an edge on some face of C_{-a} (the multiplicities of vertices are preserved). It should be noted that all arrows of the quiver of Λ_{-a} except at most two coincide with the respective arrows of the quiver of Λ_a . The arrows which don't coincide, are products of two or three arrows of the quiver of Λ_a . Again, we finish the proof only for the case when $C(a)$ is obtained from C by tilting transformation of type 3; the other cases are treated in the same way. For $i = 1, 2$ denote by b_i the edge, which precedes a_i on F_i in counter-clockwise order, i.e. b_i precedes a_i on a G -cycle (see notations in 4.2.3). Denote by μ (by ρ) the arrow in Q_e which corresponds to the angle at vertex V'_1 included between a_1 and b_1 (resp., to the angle at V'_2 included between a_2 and b_2). Define elements $\alpha_1, \beta_1, \gamma_1, \delta_1 \in \text{End}_{D^b(\Lambda)} T_a$ such that $\alpha_1\beta_1 = \mu$, $\gamma_1\delta_1 = \rho$ in $\text{End}_{D^b(\Lambda)} \bigoplus_{i \neq a} T_{ai}$. Each of these elements is induced by a morphism between two indecomposable summands of T_a :

$\alpha_1 :$

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & 0 & \longrightarrow & P_{b_1} & \longrightarrow & 0 \longrightarrow \dots \\
 & & \downarrow & & \downarrow \binom{\mu}{0} & & \downarrow \\
 \dots & \longrightarrow & 0 & \longrightarrow & P_{a_1} \oplus P_{a_2} & \xrightarrow{(\alpha, \gamma)} & P_a \longrightarrow 0 \longrightarrow \dots
 \end{array}$$

$\beta_1 :$

$$\begin{array}{ccccccc}
\cdots & \longrightarrow & 0 & \longrightarrow & P_{a_1} \oplus P_{a_2} & \xrightarrow{(\alpha, \gamma)} & P_a \longrightarrow 0 \longrightarrow \cdots \\
& & \downarrow & & \downarrow (id, 0) & & \downarrow \\
\cdots & \longrightarrow & 0 & \longrightarrow & P_{a_1} & \longrightarrow & 0 \longrightarrow \cdots
\end{array}$$

 $\gamma_1 :$

$$\begin{array}{ccccccc}
\cdots & \longrightarrow & 0 & \longrightarrow & P_{b_2} & \longrightarrow & 0 \longrightarrow \cdots \\
& & \downarrow & & \downarrow \binom{0}{\rho} & & \downarrow \\
\cdots & \longrightarrow & 0 & \longrightarrow & P_{a_1} \oplus P_{a_2} & \xrightarrow{(\alpha, \gamma)} & P_a \longrightarrow 0 \longrightarrow \cdots
\end{array}$$

 $\delta_1 :$

$$\begin{array}{ccccccc}
\cdots & \longrightarrow & 0 & \longrightarrow & P_{a_1} \oplus P_{a_2} & \xrightarrow{(\alpha, \gamma)} & P_a \longrightarrow 0 \longrightarrow \cdots \\
& & \downarrow & & \downarrow (0, id) & & \downarrow \\
\cdots & \longrightarrow & 0 & \longrightarrow & P_{a_2} & \longrightarrow & 0 \longrightarrow \cdots
\end{array}$$

The elements $\alpha_1, \beta_1, \gamma_1, \delta_1$ are not invertible, since for $i \neq a$ $H^*(T_{aa}) \neq H^*(T_{ai})$. Therefore these are the arrows μ and ρ (in Λ_{-a}) which are products of two arrows of Λ_a . Now observe that in terms of Brauer complexes, transformation of the quiver of Λ_{-a} to Λ_a is insertion of edge labeled by a , incident with V'_1 and V'_2 into the union of faces F_1 and F_2 . $\square$

Corollary 4.6. Let Λ_1 and Λ_2 be symmetric SB-algebras, let C_1 and C_2 be their Brauer complexes. Suppose that C_2 can be obtained from C_1 by a sequence of tilting transformations. Then Λ_1 and Λ_2 are derived equivalent.

Proof. The statement follows from Proposition 4.4, Lemma 4.5 and the Rickard's Theorem. $\square$

Example 4.7. Consider decagons D_1 and D_2 . Fix an orientation on each of decagons. Mark the edges of D_1 (of D_2) with letters a, b, c, d, e so that they form a word $abcdeabcde$ (resp., $abcdeadebc$) in counter-clockwise order. In each decagon, identify the edges which are marked by the same letter in such way that the resulting manifolds are oriented. It's easy to see that both complexes (we call them C_1 and C_2) have 2 vertices, 5 edges, one face, i.e. they are homeomorphic to a sphere with two handles. Moreover, the 1-skeletons of C_1 and C_2 are bipartite graphs. But these complexes cannot be obtained from each other by tilting transformations: any complex C' , obtained from the complex C_1 , is isomorphic to C_1 . This construction gives pairs of symmetric SB-algebras of genus 2, for which the methods given in present paper are not enough to determine whether they are derived equivalent or not.

5 Algebras of genus 0

Now we prove that if Brauer complex of Λ is homeomorphic to a sphere, then the multiset of perimeters of its faces and the multiset of multiplicities of vertices determine the class of derived equivalence of Λ . For a start, we don't take into consideration the multiplicities of vertices, i.e. we consider graphs with non-labeled vertices. We fix plane graphs Γ_1 and Γ_2 with the same multisets of perimeters of faces and show that Γ_2 can be obtained from Γ_1 by a sequence of tilting transformations (statements from Lemma 5.2 to Proposition 5.18).

Definition 5.1. Graphs which can be obtained from each other by a sequence of tilting transformations will be called *chain equivalent* graphs.

Lemma 5.2. Let Γ be a plane graph, $A \in V(\Gamma)$. There exists a plane graph Γ' , chain equivalent to Γ , in which the vertex A is incident with all edges and one of the following conditions holds:

1. Γ' has no loops
2. Each edge of Γ' is either a leaf or a loop at vertex A (i.e., there are no multiedges in Γ' except for loops).

Definition 5.3. Plane graph of this form is called a *reduced graph*.

Proof. Consider among graphs, which are chain equivalent to Γ , a graph Γ' with a maximal degree of A . Observe that all edges of Γ' are incident with A . Indeed, otherwise there are vertices $B, C \neq A$ and an edge $e \in E(B, C)$ such that either B or C is incident with A (without loss of generality, B) and such that the edge $\pi_B(e) \in E(A, B)$. If $B \neq C$, we apply to e a transformation of type 3. If $B = C$, we apply to e a transformation of type 2 so that e shifts from B to A . Thus the degree of A can be increased, a contradiction. It follows that there are three types of edges in Γ' :

- a) a loop at vertex A ;
- b) edges which form a multiedge incident with A ;
- c) a leaf (A, X) .

For further convenience, elements of type a) don't belong to type b). We show that in Γ' edges of types a) and b) cannot exist simultaneously. Suppose that there is a loop a , leaves $a_1 = \pi_A(a)$, $a_2 = \pi_A(a_1), \dots, a_s = \pi_A(a_{s-1})$ and an edge $b = \pi_A(a_s)$ of type b). Consider the edge $c = \pi_B(b)$. By transformations of type 1, we shift $a_1, \dots, a_s$ along a . Now there are no edges between a and b around A , and we can apply a transformation of type 3 to the edge b and b becomes a loop. This increases the degree of A , a contradiction. $\square$

Definition 5.4. A reduced graph which has no loops is called a *reduced graph of type 1*.

Observe that the border of any face of a reduced graph of type 1 is formed by several pairs of edges $(A, B_1), \dots, (A, B_k)$ and by several leaves (any leaf is counted in the perimeter of the face twice). Observe that a reduced graph of type 1 is bipartite.

Definition 5.5. A reduced graph which has loops is called a *reduced graph of type 2*.

In a reduced graph of type 2, any edge which is not a leaf is a loop. Observe that a reduced graph of type 2 is not bipartite.

Let Γ'_1 and Γ'_2 be reduced graphs, chain equivalent to Γ_1 and to Γ_2 , respectively. By Proposition 3.7, Γ'_1 and Γ'_2 are of the same type. We will show that all reduced graphs of the same type, with the same multisets of perimeters of faces, are chain equivalent:

I. Reduced graphs of type 1. Fix a graph Γ of type 1.

Lemma 5.6. Each reduced graph Γ of type 1 is chain equivalent to a reduced graph Γ' of type 1, which has at most two non-dangling vertices.

Proof. Consider among reduced graphs, which are chain equivalent to Γ , a graph Γ' with maximal number of dangling vertices. Let A be the vertex of Γ' , which is incident with all edges. We show that Γ' has at most two non-dangling vertices (including A). Indeed, let $b \in E(A, B)$ and $c \in E(A, C)$ be two edges of type 2 ($B \neq C$) such that there are only leaves between b and c in clockwise order around A . As above, by transformations of type 1 we obtain a graph, in which there are no leaves between b and c (around A). Suppose that C has degree 2. Applying the transformation of type 3 to c (shift along b), we get a reduced graph with a greater number of leaves, since C becomes a leaf. In order to transform C to a leaf when $\deg(C) = r$, we need to carry out the same operations with $r - 1$ edges, which are incident with C . $\square$

Consider a reduced graph Γ' which was obtained in lemma 5.6. It is easy to see that the faces of Γ' and the edges of Γ' which are not leaves can be cyclically numbered by $1, 2, \dots, g$ so that the border of the face number i consists of the edges number i and $i + 1$ and several inner leaves. It should be mentioned that if $g = 1$ then Γ' is a tree in a form of star, and we get Brauer trees, which were studied by Rickard in [7], as a first application of the criterion of derived equivalence.

Lemma 5.7. *Graph Γ' is chain equivalent to a graph of the same form (i.e., as in lemma 5.6), in which the perimeters of faces are in ascending ordering.*

Proof. It's enough to show how to 'transpose' two faces, see Figure 4. $\square$




Figure 4: to Lemma 5.7

We see that any bipartite plane graph is chain equivalent to a (unique) *canonical representative* (we will also say "a *graph in canonical form*") — a graph in which the perimeters of faces are in ascending ordering. Two graphs with the same multisets of perimeters are chain equivalent to the same canonical representative, and therefore they are chain equivalent to each other.

II. Reduced graphs of type 2.

Consider a reduced graph Γ of type 2. First suppose that A is the only vertex of Γ , i.e. **all edges of Γ are loops** and $n = g - 1$, where n is the number of vertices of Q_e and g is the number of G -cycles. Consider a graph $T = T(\Gamma)$, which is plane dual to Γ . T is a tree with $g - 1$ edges and g vertices. Observe that the transformations of type 1 cannot be applied to Γ . The transformations of types 2 and 3 can be described in terms of T as follows.

- Transformation of type 2. A leaf V_1V_2 of T (with dangling vertex V_1) is shifted around V_2 in arbitrary way. This transformation of a plane labeled tree will be called a *flip-over*.
- Transformation of type 3. Suppose that $\pi_{V_1}^{-1}(V_1V_2) = V_1V_3$ and that $\pi_{V_2}^{-1}(V_1V_2) = V_2V_4$. Replace edges V_1V_3 and V_2V_4 with edges V_1V_4 and V_2V_3 in a way that $\pi_{V_1}(V_1V_2) = V_1V_4$ and $\pi_{V_2}(V_1V_2) = V_2V_3$. This transformation of a plane labeled tree will be called a *flip* (see Figure 5; an arc between two edges in the pictures denotes absence of other edges).




Figure 5: Flip

Definition 5.8. Plane trees with labeled vertices, which can be obtained from each other by flips and flip-overs, are called *equivalent*. Clearly, equivalent trees are dual to chain-equivalent graphs.

Proposition 5.9. *Two plane trees with the same multisets of labeled vertices and the same degrees of correspondent vertices are equivalent.*

We need the following lemma.

Lemma 5.10. *Let V_1V_2 be a leaf in a plane tree T with dangling vertex V_1 . Let $V_1, V_2, \dots, V_r$ be a path in T such that V_r is a non-dangling vertex. Then T is equivalent to a tree, in which V_1 is adjacent with V_r .*

Proof. The proof is by induction on r . For $r = 2$ the claim is trivial. Suppose that there is a number $i \in \{1, \dots, r\}$ such that $\deg(V_i) \geq 3$. Consider the minimal such i . Without loss of generality we assume that $V_{i+1} \neq V$, where V is such vertex that $\pi_{V_i}(V_iV_{i-1}) = V_iV$. If $i \neq 2$, replace edges $V_{i-1}V_{i-2}$ and V_iV with V_iV_{i-2} and $V_{i-1}V$ by a flip. Otherwise, we make V_2V_3 follow V_2V_1 by several flip-overs, and then make the above flip. The distance between V_1 and V_r decreases, and we apply the inductive hypothesis. If i cannot be defined, consider the unique vertex $V_{r+1} \neq V_{r-1}$ adjacent with V_r . Replace $V_{r-1}V_{r-2}$ and V_rV_{r+1} with V_rV_{r-2} and $V_{r-1}V_{r+1}$ by a flip. Again, the distance between V_1 and V_r is decreased, and we apply the inductive hypothesis. $\square$

Now we prove Proposition 5.9.

Proof. The proof is by induction on the number of vertices. For $g = 1$ the claim is trivial. Let T_1 and T_2 be two plane trees with g vertices. Let a dangling vertex V be adjacent with V_1 in T_1 and with V_2 in T_2 . By Lemma 5.10, we can replace T_1 with an equivalent tree T_3 in which V is adjacent with V_2 . Let T_3^1 and T_2^1 be the trees, obtained from T_3 and T_2 by removing V with

the corresponding edge. They have the same degrees of correspondent vertices, and therefore they are equivalent by inductive hypothesis. It remains to show that it is still possible to carry out the sequence of transformations, which puts T_3^1 to T_2^1 , when edge V_2V is not deleted. After these transformation we will be able to flip-over the edge V_2V to the required place.

Start to apply the above sequence of transformations to T_3 . We can encounter difficulties in the following cases:

- When in T_3 the edge V_2V is between two subsequent edges (around V_2) of T_1^3 and doesn't allow to make a flip. We cope with this by an arbitrary flip-over of V_2V .
- If V_2 is a dangling vertex in T_3^1 , incident with an edge V_2V_3 , and in T_3^1 it is possible to make a flip-over of V_2V_3 . In T_3 instead of this flip-over we make the following sequence of transformations (Figure 6).




Figure 6: to Proposition 5.9

This finishes the proof. □

Now suppose that **there are dangling vertices in Γ** .

Definition 5.11. *External perimeter* of a face is the number of its edges, which separate it from other faces (in our case, these are loops).

Definition 5.12. *Reduction* of graph Γ is a graph $\mathcal{R}(\Gamma)$ which is obtained from Γ by removing all dangling vertices.

Proposition 5.13. *Let Γ_1 and Γ_2 be reduced graphs of type 2. Suppose that there is a tilting transformation p which puts $\mathcal{R}(\Gamma_1)$ to $\mathcal{R}(\Gamma_2)$. Suppose also that the correspondent labeled faces of Γ_1 and of Γ_2 have the same number of edges. Then graphs Γ_1 and Γ_2 are chain equivalent.*

Proof. Let l be the loop, which is shifted by p and let F_1 and F_2 be the faces separated by l . We need to obtain a sequence of transformations which would serve as an analogue of p for Γ_1 . Figure 7 illustrates the case when F_1 has inner leaves and l is the only loop on the border of F_1 . The case when there are other loops on the border of F_1 is even easier: the analogue of p is a transformation of type 3, made after necessary flip-overs of leaves. □




Figure 7: to Proposition 5.13

Remark 5.14. It follows from Propositions 5.9 and 5.13 that the class of chain equivalence of a reduced graph of type 2 is determined by the multiset of pairs $(P(F_i), p(F_i))$, where $P(F_i)$ is the perimeter and $p(F_i)$ is the external perimeter of the face F_i .

Definition 5.15. The multiset of pairs $(P(F_i), p(F_i))$ will be called a *multiset of double perimeters* of graph Γ .

Proposition 5.16. Let Γ_1 and Γ_2 be reduced graphs of type 2 with the same multisets of perimeters of faces. Then there exists a reduced graph Γ_3 of type 2, chain equivalent to Γ_1 , such that the multisets of double perimeters of Γ_2 and Γ_3 are the same.

Proof. Let $\{(P_i, p_i)\}$ be the multiset of double perimeters of Γ_2 , let $\{(P_i, p_i^1)\}$ be the multiset of double perimeters of Γ_1 , for $i = 1, \dots, g$. Observe that $p_i \equiv P_i \equiv p_i^1 \pmod{2}$ for each $i \in \{1, \dots, g\}$ and that $\sum_i p_i = 2g - 2 = \sum_i p_i^1$. Set $q_i = p_i^1$ for each i . Consider the following algorithm of 'transformation' of the multiset $\{q_i\}$ to the multiset $\{p_i\}$. Below we will show that for each step of this algorithm there is a chain equivalence of graphs, which properly changes their external perimeters.

Consider maximal k such that $q_i = p_i$ for all $i < k$.

1. If $q_k < p_k$ then $q_j > p_j$ for some $j > k$. Replace q_k with $q_k + 2$ and replace q_j with $q_j - 2$.
2. Otherwise $q_k > p_k \geq 1$ and $q_j < p_j$ for some $j > k$. In this case we replace q_k with $q_k - 2$ and replace q_j with $q_j + 2$.

Observe that at each step the number which is decreased is greater than two, so the resulting numbers are positive. Moreover, since $q_i \leq \max(p_i, p_i^1)$, at each step $q_i \leq P_i$ for all i . Clearly, the multiset of numbers q_i can be transformed to the multiset of numbers p_i by these operations. To find the chain equivalences which correspond to these operations, we need the following lemma.

Lemma 5.17. Let T be a tree, let $V_1, V_2 \in V(T)$. If V_1 and V_2 are not both dangling vertices, then there exists a tree in which the degrees of all vertices are the same and the vertices V_1 and V_2 are adjacent.

Proof. The proof is by induction on the number of vertices in T . □

We return to the proof of Proposition 5.16. We need a sequence of tilting transformations under which the multiset of external perimeters changes in accordance to the above algorithm.

Suppose that we are to change the external perimeters q_i and q_k of faces F_i and F_k , respectively, in a graph Γ . By Lemma 5.17 and Remark 5.14, Γ can be transformed to a chain equivalent graph Γ' with the same multiset of double perimeters, such that in the dual tree $T(\Gamma')$ the vertices of degrees q_i and q_k are adjacent. Without loss of generality, we are to increase q_i . In this case $q_i < P_i$ and $q_k \geq 3$. Consider faces F_1 and F_2 of Γ' which can be described in terms of dual tree $T(\Gamma')$ as follows: $F_1 = \pi_{F_k}^{-1} F_i$, $F_2 = \pi_{F_k}^{-1} F_1$ (all faces F_1 , F_2 and F_i are different, since $q_k \geq 3$). Since $q_i < P_i$, there is at least one leaf in F_i . The following sequence of transformations finishes the proof (see Figure 8; in the picture the shifts of leaves are omitted). Thus q_i is




Figure 8: to Proposition 5.16

increased by 2 and q_k is decreased by 2, which was required. $\square$

Altogether, we get

Proposition 5.18. *Two plane graphs with the same multiset of perimeters of faces are chain equivalent.*

Now we again consider graphs with labeled vertices, i.e., we return the multiplicities of vertices into consideration. In statements from Lemma 5.19 to Theorem 5.22 we prove that *if two plane Brauer graphs with the same multisets of labels of vertices are isomorphic as non-labeled graphs, then they are chain equivalent as labeled graphs*. In view of the above arguments, it's enough to prove this for reduced graphs. Moreover, in the case of bipartite graphs we may restrict ourselves to considering graphs in canonical form. Recall that the process of putting a graph to reduced form (and to canonical form, for graphs of type 1) started with choosing an *arbitrary* vertex A . Recall also that we can arbitrarily shift leaves in a face, by tilting transformation of type 1.

I. Reduced graphs of type 1. For reduced graphs of type 1, it suffices to prove the following lemmas:

Lemma 5.19. *Let Γ be a graph in canonical form, let $B \neq A$ be the second non-dangling vertex of Γ , let F be a face. Then Γ is chain equivalent to a graph in canonical form, in which*

1. B is a dangling vertex in the face F .
2. Some vertex $C \neq A$ which belongs in Γ to F is a non-dangling vertex.
3. The other dangling vertices belong in Γ and in Γ_1 to the same faces.

Lemma 5.20. *Let Γ be a graph in canonical form, let $B \neq A$ be the second non-dangling vertex of Γ , let faces F_1 and F_2 be adjacent. Then Γ is chain equivalent to a graph in canonical form Γ_1 , in which*

1. *There is a dangling vertex which belongs in Γ to F_1 and belongs in Γ_1 to F_1 , and there is another dangling vertex which belongs in Γ to F_2 and in Γ_1 to F_2 .*
2. *The other dangling vertices belong in Γ and in Γ_1 to the same faces.*

For the proof of Lemma 5.19 see Figure 9. For the proof of Lemma 5.20 see Figure 10.




Figure 9: Proof of Lemma 5.19




Figure 10: Proof of Lemma 5.20

II. Reduced graphs of type 2. Since the dangling vertices in a face can be shifted in arbitrary way, it's enough to show how to interchange dangling vertices belonging to different faces (say, to F_1 and F_2). First consider the case when the external perimeter of F_1 or F_2 is greater than 1. Then by Lemma 5.17 and Remark 5.14, there is a sequence of tilting transformations making F_1 and F_2 adjacent. Moreover, this sequence preserves the faces to which belong the dangling vertices (see Figure 7). Therefore, in this case it's enough to show how to interchange dangling vertices which belong to adjacent faces: see Figure 11. Now consider the case when the dangling vertices which we want to interchange belong to faces, which correspond to dangling vertices of the dual tree.

Lemma 5.21. *Let T be a plane tree with labeled vertices, let V_1 and V_2 be dangling vertices of T . Suppose that T is not a chain. Then T is equivalent to a tree, in which the edges which are incident with V_1 and V_2 are incident to a common vertex V . Moreover, $\pi_V(VV_1) = VV_2$.*




Figure 11: Interchange between adjacent faces

Proof. By Remark 5.14 it is enough to find a tree with the same multiset of degrees as in T , in which some two leaves are adjacent to a common vertex. Denote the degrees of T by $r_1, \dots, r_g$ in such way that $r_1 = r_2 = 1$, $r_3 \geq 3$. Observe that the sum of numbers $d_3 - 2, d_4, \dots, d_g$ equals $2g - 6$. It can be shown by induction on g that there is a tree T' , in which these numbers are the degrees of vertices. To obtain the needed tree, we add two leaves to the vertex of T' of degree $r_3 - 2$. $\square$




Figure 12: Interchange between "dangling" faces

We see that if T is not a chain, then it suffices to show how to interchange dangling vertices between two "dangling" faces, which have a common adjacent face: see Figure 12. It remains to examine the case when $T(\Gamma)$ is a chain, and F_1 and F_2 correspond to the two dangling vertices of $T(\Gamma)$. If some other face of Γ contains a dangling vertex, the needed interchange comes to three interchanges of the above form. In Figure 13 is shown how to interchange leafs in case when the rest faces have perimeter 2. (For the graph in the picture $g = 4$, and this case fully represents the general case.)

This finishes the proof of the main theorem in this section:

Theorem 5.22. *Let Λ_1 and Λ_2 be symmetric SB-algebras of genus 0. Then Λ_1 and Λ_2 are derived equivalent if and only if their Brauer complexes have the same multisets of perimeters of faces and the same multisets of labels on vertices.*




Figure 13: Chain case

References

- [1] M. A. Antipov, A. I. Generalov, *Yoneda algebras of symmetric special biserial algebras are finitely generated*, Algebra and Analysis 2005, Vol. 17, pp. 1–23 (in Russian).
- [2] M. A. Antipov, *Invariants of stable equivalence of symmetric special biserial algebras*, Zapiski Nauchnykh Seminarov POMI, Vol. 330, 2006, pp. 5–28 (in Russian).
- [3] Th. Holm, *Derived equivalence classification of algebras of dihedral, semidihedral, and quaternion type*, J. Algebra. 1999, Vol. 211, 1, pp. 159–205.
- [4] S. K. Lando, A. K. Zvonkin, *Graphs on Surfaces and their Applications*, 2003
- [5] Z. Pogorzaly, *On a construction of algebras stably equivalent to selfinjective special biserial algebras*, Ann. Sci. Math. Quebec 1993, Vol. 17, 1, pp. 65–97.
- [6] J. Rickard, *Morita theory for derive categories*, J.Lonon Math. Soc., 1989, Vol. 39, pp. 436–456.
- [7] J. Rickard, *Derived categories and stable equivalence*, Jouranal Pure Appl. Algebra, 1989, Vol. 61, pp. 303–317.